

R-Parity violating minimal supergravity at the LHC

B. C. Allanach^{1 a}, M. A. Bernhardt^{2 b}, H. K. Dreiner^{2 c}, S. Grab^{2 d}, C. H. Kom^{1 e}, and P. Richardson^{3 f}

¹ DAMTP, University of Cambridge, Cambridge, UK

² Physikalisches Institut, University of Bonn, Bonn, Germany

³ IPPP, University of Durham, Durham, UK

Abstract. We consider the case where supersymmetry with broken R-parity is embedded in the minimal supergravity model (mSUGRA). This alters the standard mSUGRA spectrum and opens a wide range in parameter space, where the scalar tau is the lightest supersymmetric particle, instead of the lightest neutralino. We study the resulting LHC phenomenology. Promising signatures would be detached vertices from long-lived staus, multi lepton final states and multi-tau final states. We investigate in detail the corresponding cross sections and decay rates in characteristic benchmark scenarios.

PACS. 04.65.+e Supergravity – 12.60.Jv Supersymmetric models – 14.80.Ly Supersymmetric partners of known particles

1 R-Parity violating minimal supergravity model

The general renormalizable superpotential of the minimal supersymmetric extension of the SM includes lepton and baryon number violating terms [1]:

$$W_{R_p} = \epsilon_{ab} \left[\frac{1}{2} \lambda_{ijk} L_i^a L_j^b \bar{E}_k + \lambda'_{ijk} L_i^a L_j^b \bar{D}_k \right] + \epsilon_{ab} \kappa^i L_i^a H_2^b + \frac{1}{2} \epsilon_{xyz} \lambda''_{ijk} \bar{U}_i^x \bar{D}_j^y \bar{D}_k^z. \quad (1)$$

The lepton and baryon number violating interactions in Eq. (1) lead to rapid proton decay [2], if simultaneously present. Therefore the supersymmetric Lagrangian must respect an additional discrete symmetry. The most widely studied scenario is R-Parity [3], R_p , for which $W_{R_p} = 0$. This scenario is conventionally named the minimal supersymmetric standard model (MSSM). A detailed analysis of all discrete symmetries within the minimal supersymmetric extension of the SM can be found in Ref. [4].

The MSSM with R_p has 124 free parameters [5], and without R_p more than 200. Thus it is mandatory to consider simpler models with well motivated boundary conditions at the unification scale, M_{GUT} , because such an extensive parameter space is intractable for a systematic phenomenological analyses at colliders.




Fig. 1. Parameter space around SPS1a, with $A_0 = -100$ GeV, $\tan\beta = 10$, and $\text{sgn}(\mu)=+1$. The bar on the right displays the mass of the lightest stau $m_{\tilde{\tau}}$. On the left side, we show in black the excluded region due to tachyons and LEP2 Higgs exclusion bounds. The black contour distinguishes between areas with $\tilde{\tau}$ -LSP and $\tilde{\chi}_1^0$ -LSP [9].

The most widely studied model is mSUGRA [6] with radiative electroweak symmetry breaking [7] and conserved R_p . This model has only five parameters (defined at M_{GUT})

$$M_0, \quad M_{1/2}, \quad A_0, \quad \tan\beta, \quad \text{sgn}(\mu). \quad (2)$$

Here M_0 is the universal soft-breaking scalar mass, $M_{1/2}$ denotes the universal gaugino mass, A_0 is the universal soft-breaking scalar interaction and $\tan\beta$ is the ratio of the vacuum expectation values of the two Higgs doublets. The sign of the Higgs mixing parameter is $\text{sgn}(\mu)$.

^a Email: b.c.allanach@damtp.cam.ac.uk

^b Email: markus@th.physik.uni-bonn.de

^c Email: dreiner@th.physik.uni-bonn.de

^d Speaker, Email: sgrab@th.physik.uni-bonn.de

^e Email: c.kom@damtp.cam.ac.uk

^f Email: Peter.Richardson@durham.ac.uk

$$\begin{array}{ll}
\tilde{\tau}^- \rightarrow \tau^- \bar{\nu}_\mu \bar{d}d & \tilde{\tau}^- \rightarrow \tau^- \nu_\mu d\bar{d} \\
\tilde{\tau}^- \rightarrow \tau^- \mu^+ \bar{u}d & \tilde{\tau}^- \rightarrow \tau^- \mu^- u\bar{d} \\
\tilde{\tau}^- \rightarrow \nu_\tau \mu^- u\bar{u} & \tilde{\tau}^- \rightarrow \nu_\tau \nu_\mu d\bar{u} \\
\tilde{\tau}^- \rightarrow \nu_\tau \mu^- d\bar{d} & \tilde{\tau}^- \rightarrow \nu_\tau \bar{\nu}_\mu \bar{u}d
\end{array}$$

Figure 1: Mass spectrum of the dark sector. The plot shows the mass m in GeV on the y-axis (0 to 1000) versus an unlabeled x-axis. Various particles are plotted as horizontal lines with labels. Red lines represent h^0 , $h^\pm$, and A^0, H^0 . Black lines represent $\tilde{l}_L^\pm$ and $\tilde{l}_R^\pm$. Blue lines represent various fermions and scalars: $\tilde{q}_L^\pm$, $\tilde{q}_R^\pm$, $\tilde{b}_1^\pm$, $\tilde{b}_2^\pm$, $\tilde{g}$, $\tilde{l}_1^\pm$, $\tilde{l}_2^\pm$, and various χ particles. The mass values are approximately: h^0 at 125 GeV, $h^\pm$ at 125 GeV, A^0, H^0 at 560 GeV, $\tilde{l}_L^\pm$ at 250 GeV, $\tilde{l}_R^\pm$ at 250 GeV, $\tilde{q}_L^\pm$ at 850 GeV, $\tilde{q}_R^\pm$ at 850 GeV, $\tilde{b}_1^\pm$ at 850 GeV, $\tilde{b}_2^\pm$ at 850 GeV, $\tilde{g}$ at 950 GeV, $\tilde{l}_1^\pm$ at 650 GeV, $\tilde{l}_2^\pm$ at 850 GeV, and various χ particles at 500 GeV.

- For $\Lambda \gtrsim O(10^{-2})$, the $\mathcal{R}_p$ operator can significantly change the decay chains of the sparticles. Also the mass spectrum can be significantly affected [9]. Furthermore single sparticle production may dominate, e.g., single slepton production for $\lambda'_{ijk} \neq 0$ [13].

Table 2. Signal rates for benchmark scenario BC1. The numbers of electrons or muons of positive charge (first row) and negative charge (second row) are shown. We also give the numbers of taus with positive (third row) and negative charge (fourth row). Furthermore, we mention, if there is missing p_T (neglecting that from τ decay) due to neutrinos (fifth row) and we present the probability of the different signatures (sixth row). Each event is accompanied by 2–4 jets. $\sigma_{tot} = 4.8 \times 10^3$ fb.

e^+ or μ^+	e^- or μ^-	τ^+	τ^-	$\cancel{p}_T$	event fraction
2	2	2	2	yes	35 %
3	2	2	2	yes	12 %
2	3	2	2	yes	8.3 %
3	3	2	2	yes	7.3 %
2	2	2	1	yes	4.7 %
2	2	3	2	yes	4.3 %
2	2	3	3	yes	1.4 %
4	3	2	2	yes	1.1 %

3 Stau LSP phenomenology at the LHC.

3.1 Benchmark scenarios

In Ref. [9], four benchmark scenarios, BC1–BC4, within the framework of $\tilde{R}_p$ mSUGRA were proposed. Three scenarios, BC1, BC2, BC4, contain a $\tilde{\tau}$ -LSP and BC3 a sneutrino-LSP. In the following we will consider two of these scenarios, namely:

- **BC1:** $M_0 = A_0 = 0$ GeV, $M_{1/2} = 400$ GeV, $\tan \beta = 13$, $\text{sgn}(\mu) = +1$, $\lambda_{121} = 0.032$ at M_{GUT} .
- **BC2:** $M_0 = A_0 = 0$ GeV, $M_{1/2} = 400$ GeV, $\tan \beta = 13$, $\text{sgn}(\mu) = +1$, $\lambda'_{311} = 3.5 \times 10^{-7}$ at M_{GUT} .

Both scenarios differ only by the non-vanishing $\tilde{R}_p$ operator. The mass spectrum is shown in Fig. 3. The $\tilde{\tau}$ is the LSP with $m_{\tilde{\tau}} = 148$ GeV. The spectrum is calculated with an $\tilde{R}_p$ version of **SOFTSUSY** [16]. The effect of the $\tilde{R}_p$ operators on the sparticle masses is smaller than 1 GeV and thus negligible. Note that for BC1 and BC2, the NLSP to NNNLSP are nearly degenerate in mass ($m_{\tilde{e}_R} = 161$ GeV, $m_{\tilde{\mu}_R} = 161$ GeV, $m_{\tilde{\chi}_1^0} = 162$ GeV).

3.2 Signal rates at the LHC

We now investigate the signatures at the LHC at parton level. We use a modified version of **HERWIG** [15], which contains all 2- and 4-body decays of the $\tilde{\tau}$ -LSP. All other decay rates of sparticles are calculated with **ISAWIG1.200** and **ISAJET7.64** [14].

We show the signal rates of the benchmark scenario BC1 in Table 2. The total cross section for all sparticle pair production processes is $\sigma_{tot} = 4.8 \times 10^3$ fb. Each event is accompanied by four or more electrons or muons in the final state. These leptons allow for triggering and for reconstruction of the event. Most of these leptons originate from the 4-body decays of the $\tilde{\tau}$ -LSP via λ_{121} , see Table 3. We also observe in

Table 3. Dominant branching ratios (BR) of the LSP, $\dots$, NNNLSP for benchmark scenario BC1. $\tilde{R}_p$ BRs are bold face [9].

	mass [GeV]	channel	BR	channel	BR
$\tilde{\tau}_1^-$	148	$\mu^+ \bar{\nu}_e e^- \tau^-$	32 %	$e^+ \bar{\nu}_\mu e^- \tau^-$	32 %
		$\mu^- \nu_e e^+ \tau^-$	18 %	$e^- \nu_\mu e^+ \tau^-$	18 %
$\tilde{e}_R^-$	161	$e^- \nu_\mu$	50 %	$\mu^- \nu_e$	50 %
$\tilde{\mu}_R^-$	161	$\tilde{\tau}^+ \mu^- \tau^-$	51 %	$\tilde{\tau}^- \mu^- \tau^+$	49 %
$\tilde{\chi}_1^0$	162	$\tilde{\tau}_1^+ \tau^-$	50 %	$\tilde{\tau}_1^- \tau^+$	50 %

Table 4. Same as Table 2, but for benchmark scenario BC2. Each event is accompanied by 6–8 jets.

e^+ or μ^+	e^- or μ^-	τ^+	τ^-	$\cancel{p}_T$	event fraction
0	0	1	1	no	14 %
0	0	2	0	no	7.1 %
0	0	0	2	no	6.8 %
1	0	1	1	yes	6.5 %
0	0	1	1	yes	4.5 %
1	0	0	2	yes	3.3 %
1	0	2	0	yes	3.2 %
1	1	1	1	yes	2.4 %

Table 5. Same as Tab. 3, but for benchmark scenario BC2 [9].

	mass [GeV]	channel	BR	channel	BR
$\tilde{\tau}_1^-$	148	$\bar{u}d$	100 %		
$\tilde{e}_R^-$	161	$\tilde{\tau}_1^+ e^- \tau^-$	51 %	$\tilde{\tau}_1^- e^- \tau^+$	49 %
$\tilde{\mu}_R^-$	161	$\tilde{\tau}_1^+ \mu^- \tau^-$	51 %	$\tilde{\tau}_1^- \mu^- \tau^+$	49 %
$\tilde{\chi}_1^0$	162	$\tilde{\tau}_1^+ \tau^-$	50 %	$\tilde{\tau}_1^- \tau^+$	50 %

Table 3, that the $\tilde{e}_R$ decays into a lepton and neutrino, since it directly couples to the $L_1 L_2 \bar{E}_1$ operator. Each 4-body decay and each $\tilde{\chi}_1^0$ decay is accompanied by a tau, which leads to 4 taus in most of the events. Thus a good tau identification would help to identify this scenario. The low- p_T taus will be invisible, yet the high energy tail, $p_T \geq 30$ GeV, see Fig. 4, is useful. The small number of jets (2–4 jets) give only a small combinatorial background for tau identification via hadronic decays. Finally, each 4-body decay includes a final-state neutrino, resulting in missing transverse momentum, $\cancel{p}_T$, shown in Fig. 4. This is however reduced compared to the R_p -MSSM, where the $\cancel{p}_T$ peaks roughly around 100 GeV.

We now move on to BC2, where we have $\lambda'_{311} = 3.5 \times 10^{-7}$ at M_{GUT} . This changes the LHC signatures in a significant way, as can be seen in Table 4. Most of the electrons and muons are absent. The $\tilde{\tau}$ -LSP is completely dominated by 2-body decays into two jets, see Table 5. Unlike in BC1, the LSP decays are neutrinoless, meaning that SUSY events do not necessarily have the classical signature of missing transverse momentum. Most events are accompanied by two taus, which originate mainly from $\tilde{\chi}_1^0$ decays, see Table 5. These taus are much harder to identify compared to BC1. On the one hand, the average p_T values of the




Fig. 4. p_T distribution of the τ from $\tilde{\tau} \rightarrow \tau + X$ (left) and p_T distribution of the neutrinos (right) in benchmark scenario BC1.




Fig. 5. p_T distribution of the τ from $\tilde{\chi}_1^0 \rightarrow \tau \tilde{\tau}$ (left) and p_T distribution of the d-jets from $\tilde{\tau} \rightarrow u d$ (right) in benchmark scenario BC2.

taus are smaller, as can be seen in Fig. 5. On the other hand, the additional 6-8 jets in each event makes their identification difficult. But there will be a detached vertex, since the decay length in the rest frame of the $\tilde{\tau}$ -LSP is $c\tau_{\tilde{\tau}} \approx 0.3$ mm. The two jets from the 2-body decay make it possible to reconstruct the $\tilde{\tau}$ -LSP mass. We show the p_T distribution of one of these jets in Fig. 5.

4 Summary and conclusion

We have shown that minimal supergravity (mSUGRA) models often provide a scalar tau instead of the lightest neutralino as the lightest supersymmetric particle (LSP). If R-Parity is violated, the stau will decay into SM particles and does not contribute to dark matter. This reopens a large part of the mSUGRA parameter space, where the stau is the LSP. We have analysed in detail the LHC phenomenology of two benchmark scenarios with a stau-LSP. Although both scenarios possess the same mass spectrum, their signatures are different. In the first benchmark scenario, BC1, the stau-LSP decays via a 4-body decay into a tau, two leptons and one neutrino. In the second benchmark scenario, BC2, the stau-LSP decays into two jets. We have to be prepared to observe SUSY events at the

LHC, which are different from the standard signatures. In particular SUSY events must not have large missing transverse momentum.

We conclude that further investigations of stau-LSP models from the phenomenological as well as from the experimental side are required. We have to classify and analyse all possibilities of the R-Parity violating mSUGRA parameter space, if we want to discover supersymmetry and its connection to physics at the GUT scale.

5 Acknowledgments

The authors want to thank Klaus Desch and Sebastian Fleischmann for helpful discussions. MB thanks DAMTP, Cambridge, and IPPP, Durham, for repeatedly offered, warm hospitality. SG thanks the Deutsche Telekom Stiftung for financial support. This work has been partially supported by STFC.

References

1. H. K. Dreiner, arXiv:hep-ph/9707435.
2. J. L. Goity and M. Sher, Phys. Lett. B **346** (1995) 69 [Erratum-ibid. B **385** (1996) 500]; A. Y. Smirnov and F. Vissani, Phys. Lett. B **380** (1996) 317; A. Y. Smirnov and F. Vissani, Nucl. Phys. B **460** (1996) 37.
3. G. Farrar and P. Fayet, Phys. Lett. B **76** (1978) 575.
4. L. E. Ibanez and G. G. Ross, Nucl. Phys. B **368** (1992) 3; H. K. Dreiner, C. Luhn and M. Thormeier, Phys. Rev. D **73**, (2006) 075007.
5. H. E. Haber, Nucl. Phys. Proc. Suppl. **62** (1998) 469.
6. A. Chamseddine, R. Arnowitt and P. Nath, Phys. Rev. Lett. **49** (1982) 970; L. Alvarez-Gaume, M. Claudson and M. Wise, Nucl. Phys. B **207** (1982) 96; L. Ibanez, Phys. Lett. B **118** (1982) 73. S. Soni and H. Weldon, Phys. Lett. B **126** (1983) 215. L. Hall, J. Lykken and S. Weinberg, Phys. Rev. D **27** (1983) 2359.
7. L. Ibanez and G. Ross, Phys. Lett. B **110** (1982) 215.
8. B. C. Allanach *et al.*, arXiv:hep-ph/0202233.
9. B. C. Allanach, M. A. Bernhardt, H. K. Dreiner, C. H. Kom and P. Richardson, Phys. Rev. D **75** (2007) 035002.
10. J. Ellis, *et al.*, Nucl. Phys. B **238** (1984) 453.
11. W. Buchmuller, L. Covi, K. Hamaguchi, A. Ibarra and T. Yanagida, JHEP **0703** (2007) 037.
12. B. C. Allanach, A. Dedes and H. K. Dreiner, Phys. Rev. D **69** (2004) 115002 [Erratum-ibid. D **72** (2005) 079902].
13. S. Dimopoulos and L. J. Hall, Phys. Lett. B **207** (1988) 210; S. Dimopoulos, R. Esmailzadeh, L. J. Hall and G. D. Starkman, Phys. Rev. D **41** (1990) 2099; H. K. Dreiner, S. Grab, M. Kramer and M. K. Trenkel, Phys. Rev. D **75** (2007) 035003.
14. F. E. Paige, S. D. Protopopescu, H. Baer and X. Tata, arXiv:hep-ph/0312045.
15. G. Corcella *et al.*, JHEP **0101** (2001) 010; S. Moretti, K. Odagiri, P. Richardson, M.H. Seymour and B.R. Webber, JHEP **0204** (2002) 028.
16. B. C. Allanach, M. A. Bernhardt, preprint: BONN-TH-2007-06.